

This is the title of my talk.

And the subtitle of my talk.

My Full Name

The Name Of My Research Group

The Place Where I Work

Where I'm Speaking, When I'm Speaking



**Your
Logo goes
Here**

arXiv:xxxx.xxxx

arXiv:yyyy.yyyy

arXiv:zzzz.zzzz

This sample presentation uses lots of different colors and background gradients. The point is to show what is possible. Be conservative with your choices. Use patterns sparingly, and use color when it emphasizes or clarifies what you are trying to explain.

Outline

Introduction

First Section

Final Remarks

Macros for Boxes

Backgrounds

Custom Colors

Outline

Introduction

First Section

Final Remarks

Macros for Boxes

Backgrounds

Custom Colors

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

- First Reason.

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

- First Reason.
- Second Reason.

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

- First Reason.
- Second Reason.
- Third Reason.

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

- First Reason.
- Second Reason.
- Third Reason.

This is in a red box, so people will assume that it is important!

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

- First Reason.
- Second Reason.
- Third Reason.

This is in a red box, so people will assume that it is important!

You might also be interested in the following points

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

- First Reason.
- Second Reason.
- Third Reason.

This is in a red box, so people will assume that it is important!

You might also be interested in the following points

1. Point One.

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

- First Reason.
- Second Reason.
- Third Reason.

This is in a red box, so people will assume that it is important!

You might also be interested in the following points

1. Point One.
2. Point Two.

Why Am I Here?

(And why should you care?)

I am here giving a talk, so I must have something important to say. Let me try to convince you of this by providing a list of reasons. They will appear one at a time.

- First Reason.
- Second Reason.
- Third Reason.

This is in a red box, so people will assume that it is important!

You might also be interested in the following points

1. Point One.
2. Point Two.
3. Point Three.

The Semiclassical Limit

Consider the Euclidean path integral

$$\mathcal{Z} = \int \mathcal{D}g \mathcal{D}X \exp \left(-\frac{1}{\hbar} I_E[g, X] \right) .$$

The Semiclassical Limit

Consider the Euclidean path integral

$$\mathcal{Z} = \int \mathcal{D}g \mathcal{D}X \exp \left(-\frac{1}{\hbar} I_E[g, X] \right) .$$

- In the semiclassical ($\hbar \rightarrow 0$) limit it should be dominated by contributions from classical solutions.

The Semiclassical Limit

Consider the Euclidean path integral

$$\mathcal{Z} = \int \mathcal{D}g \mathcal{D}X \exp \left(-\frac{1}{\hbar} I_E[g, X] \right) .$$

- In the semiclassical ($\hbar \rightarrow 0$) limit it should be dominated by contributions from classical solutions.
- We want to explore the relationship between \mathcal{Z} and the Euclidean partition function

$$\mathcal{Z} \sim e^{-\beta \Omega}$$

The Semiclassical Limit

Consider the Euclidean path integral

$$\mathcal{Z} = \int \mathcal{D}g \mathcal{D}X \exp \left(-\frac{1}{\hbar} I_E[g, X] \right) .$$

- In the semiclassical ($\hbar \rightarrow 0$) limit it should be dominated by contributions from classical solutions.
- We want to explore the relationship between \mathcal{Z} and the Euclidean partition function

$$\mathcal{Z} \sim e^{-\beta \Omega}$$

- Ω is the thermodynamic potential and β is the periodicity of the Euclidean time.

The Semiclassical Limit

Consider the Euclidean path integral

$$\mathcal{Z} = \int \mathcal{D}g \mathcal{D}X \exp \left(-\frac{1}{\hbar} I_E[g, X] \right) .$$

- In the semiclassical ($\hbar \rightarrow 0$) limit it should be dominated by contributions from classical solutions.
- We want to explore the relationship between \mathcal{Z} and the Euclidean partition function

$$\mathcal{Z} \sim e^{-\beta \Omega}$$

- Ω is the thermodynamic potential and β is the periodicity of the Euclidean time.
- The ensemble depends on the boundary conditions.

Saddle-point Approximation

A small perturbation around a classical solution

$$\begin{aligned} I_E[g_{cl} + \delta g, X_{cl} + \delta X] = & I_E[g_{cl}, X_{cl}] + \delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] \\ & + \frac{1}{2} \delta^2 I_E[g_{cl}, X_{cl}; \delta g, \delta X] + \dots \end{aligned}$$

Saddle-point Approximation

A small perturbation around a classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] \\ + \frac{1}{2} \delta^2 I_E[g_{cl}, X_{cl}; \delta g, \delta X] + \dots$$

- The **leading term** is the ‘on-shell’ action.

Saddle-point Approximation

A small perturbation around a classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] \\ + \frac{1}{2} \delta^2 I_E[g_{cl}, X_{cl}; \delta g, \delta X] + \dots$$

- The **leading term** is the ‘on-shell’ action.
- The **linear term** should vanish on solutions g_{cl} and X_{cl} .

Saddle-point Approximation

A small perturbation around a classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] \\ + \frac{1}{2} \delta^2 I_E[g_{cl}, X_{cl}; \delta g, \delta X] + \dots$$

- The **leading term** is the ‘on-shell’ action.
- The **linear term** should vanish on solutions g_{cl} and X_{cl} .
- Really, the linear term should *vanish*.

Saddle-point Approximation

A small perturbation around a classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] \\ + \frac{1}{2} \delta^2 I_E[g_{cl}, X_{cl}; \delta g, \delta X] + \dots$$

- The **leading term** is the ‘on-shell’ action.
- The **linear term** should vanish on solutions g_{cl} and X_{cl} .
- Really, the linear term should *vanish*.
- The **quadratic term** represents the first corrections.

Saddle-point Approximation

A small perturbation around a classical solution

$$I_E[g_{cl} + \delta g, X_{cl} + \delta X] = I_E[g_{cl}, X_{cl}] + \delta I_E[g_{cl}, X_{cl}; \delta g, \delta X] \\ + \frac{1}{2} \delta^2 I_E[g_{cl}, X_{cl}; \delta g, \delta X] + \dots$$

- The **leading term** is the ‘on-shell’ action.
- The **linear term** should vanish on solutions g_{cl} and X_{cl} .
- Really, the linear term should *vanish*.
- The **quadratic term** represents the first corrections.

$$\mathcal{Z} \sim \exp \left(-\frac{1}{\hbar} I_E[g_{cl}, X_{cl}] \right) \int \mathcal{D}\delta g \mathcal{D}\delta X \exp \left(-\frac{1}{2\hbar} \delta^2 I_E \right) \times \dots$$

The Problem

In theories with gravity, the action we use to obtain the equations of motion usually does not give a sensible path integral formulation of the theory. Typically, we encounter three problems:

The Problem

In theories with gravity, the action we use to obtain the equations of motion usually does not give a sensible path integral formulation of the theory. Typically, we encounter three problems:

1. The action diverges on-shell.

The Problem

In theories with gravity, the action we use to obtain the equations of motion usually does not give a sensible path integral formulation of the theory. Typically, we encounter three problems:

1. The action diverges on-shell.
2. The first variation of the action does not vanish for all field configurations that contribute to the path integral.

The Problem

In theories with gravity, the action we use to obtain the equations of motion usually does not give a sensible path integral formulation of the theory. Typically, we encounter three problems:

1. The action diverges on-shell.
2. The first variation of the action does not vanish for all field configurations that contribute to the path integral.

Boundary terms:

$$\delta I|_{\text{on-shell}} \sim \int_{\partial\mathcal{M}} dx \sqrt{\gamma} \left[\pi^{ab} \delta\gamma_{ab} + \pi_X \delta X \right] \neq 0$$

The Problem

In theories with gravity, the action we use to obtain the equations of motion usually does not give a sensible path integral formulation of the theory. Typically, we encounter three problems:

1. The action diverges on-shell.
2. The first variation of the action does not vanish for all field configurations that contribute to the path integral.

Boundary terms:

$$\delta I|_{\text{on-shell}} \sim \int_{\partial\mathcal{M}} dx \sqrt{\gamma} \left[\pi^{ab} \delta\gamma_{ab} + \pi_X \delta X \right] \neq 0$$

3. The Gaussian integral in the saddle point approximation may diverge.

The Problem

In theories with gravity, the action we use to obtain the equations of motion usually does not give a sensible path integral formulation of the theory. Typically, we encounter three problems:

1. The action diverges on-shell.
2. The first variation of the action does not vanish for all field configurations that contribute to the path integral.

Boundary terms:

$$\delta I|_{\text{on-shell}} \sim \int_{\partial\mathcal{M}} dx \sqrt{\gamma} \left[\pi^{ab} \delta\gamma_{ab} + \pi_X \delta X \right] \neq 0$$

3. The Gaussian integral in the saddle point approximation may diverge.

The semiclassical analysis is more involved than we might have guessed!

Outline

Introduction

First Section

Final Remarks

Macros for Boxes

Backgrounds

Custom Colors

The Einstein-Hilbert action with GHY boundary term

$$I_E = -\frac{1}{16\pi G} \int_{\mathcal{M}} d^{n+1}x \sqrt{g} (R - 2\Lambda) - \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} K$$

The Einstein-Hilbert action with GHY boundary term

$$I_E = -\frac{1}{16\pi G} \int_{\mathcal{M}} d^{n+1}x \sqrt{g} (R - 2\Lambda) - \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} K$$

- Newton's constant

The Einstein-Hilbert action with GHY boundary term

$$I_E = -\frac{1}{16\pi G} \int_{\mathcal{M}} d^{n+1}x \sqrt{g} (R - 2\Lambda) - \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} K$$

- Newton's constant
- The Ricci scalar

The Einstein-Hilbert action with GHY boundary term

$$I_E = -\frac{1}{16\pi G} \int_{\mathcal{M}} d^{n+1}x \sqrt{g} (R - 2\Lambda) - \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} K$$

- Newton's constant
- The Ricci scalar
- The Cosmological constant

The Einstein-Hilbert action with GHY boundary term

$$I_E = -\frac{1}{16\pi G} \int_{\mathcal{M}} d^{n+1}x \sqrt{g} (R - 2\Lambda) - \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} K$$

- Newton's constant
- The Ricci scalar
- The Cosmological constant
- The Extrinsic Curvature

Outline

Introduction

First Section

Final Remarks

Macros for Boxes

Backgrounds

Custom Colors

Now I will conclude my talk.

Now I will conclude my talk.

- Closing Point One

Now I will conclude my talk.

- Closing Point One
- Closing Point Two

Now I will conclude my talk.

- Closing Point One
- Closing Point Two
- Closing Point Three

Now I will conclude my talk.

- Closing Point One
- Closing Point Two
- Closing Point Three
- Much more . . .

Now I will conclude my talk.

- Closing Point One
- Closing Point Two
- Closing Point Three
- Much more . . .

Next Stage

Now I will conclude my talk.

- Closing Point One
- Closing Point Two
- Closing Point Three
- Much more . . .

Next Stage

- Future plans one

Now I will conclude my talk.

- Closing Point One
- Closing Point Two
- Closing Point Three
- Much more . . .

Next Stage

- Future plans one
- Future plans two

Now I will conclude my talk.

- Closing Point One
- Closing Point Two
- Closing Point Three
- Much more . . .

Next Stage

- Future plans one
- Future plans two
- Future plans three

Now I will conclude my talk.

- Closing Point One
- Closing Point Two
- Closing Point Three
- Much more . . .

Next Stage

- Future plans one
- Future plans two
- Future plans three

I have taught you many important things!

The next three sections explain some of the custom commands, like colored boxes and background gradients.

Outline

Introduction

First Section

Final Remarks

Macros for Boxes

Backgrounds

Custom Colors

This is an 'equation box'. You can put anything inside it. For instance, here are some equations:

$$\alpha = \beta + \gamma$$

$$\delta = \epsilon + \eta$$

Here is another example.

You can control the width of this box.
It does **not** auto-adjust.

$$\alpha = \beta + \gamma$$

$$\delta = \epsilon + \eta$$

The ‘Equation Box’ Command

The equation box macro takes two arguments

$$\backslash eqnbox\{<width>\}\{<content>\}$$

Some important points:

1. You **must** specify the width of the box ‘by hand’. This is the first argument given to the macro.
2. You can use any environment inside an equation box. This is the second argument given to the macro. It is executed inside a minipage which is then drawn inside the colored box.
3. The colors used by `\eqnbox` are fixed. If you want different colors use the `\custombox` command.

A Title

This is a 'title box'. It is like an equation box, but it has a title.

$$\alpha = \beta + \gamma$$

$$\delta = \epsilon + \eta$$

A Centered Title!

This is a 'centered title box'.

$$\alpha = \beta + \gamma$$

$$\delta = \epsilon + \eta$$

The ‘Title Box’ Command

The title boxes macros take three arguments

$$\text{\titlebox{<width>}{<title>}{<content>}}$$
$$\text{\ctitlebox{<width>}{<title>}{<content>}}$$

Some important points:

1. The same caveats that apply to `\eqnbox` also apply here.
2. Obviously, `\ctitlebox` centers the title at the top of the box.

The Custom Box

The custom box lets you define the outer and inner colors, the width of the outer line, the inner padding, and the width of the minipage inside the box.

$$I_E = -\frac{1}{16\pi G_2} \int_{\mathcal{M}} d^2x \sqrt{g} [X R - U(X) (\nabla X)^2 - 2V(X)] \\ - \frac{1}{8\pi G_2} \int_{\partial\mathcal{M}} dx \sqrt{\gamma} X K$$

Hello!

I'm an orange box
with a red outline.

The ‘Custom Box’ Command

The Custom Box macro takes six arguments

```
\custombox{<outercolor>}{<linewidth>}{<innercolor>}  
          {<inner padding>}{<width>}{<content>}
```

1. `outercolor` is the color used to draw the outer line.
2. `linewidth` is the width of the outer line. It must be given with units; i.e. 1pt or .5pt.
3. `innercolor` is the color used to fill the box.
4. `inner padding` is the space between the edge of the box and the content. Also given with units.
5. `width` is the width of the box.
6. `content` is the content. You can place anything inside a `\custombox`.

Outline

Introduction

First Section

Final Remarks

Macros for Boxes

Backgrounds

Custom Colors

Background Gradients

You can place a gradient in the background using the 'bggradient' command. It takes two arguments: the top color and the bottom color.

```
\bggradient{<top color>}{<bottom color>}
```

For example, this page was produced with

```
\bggradient{orange!20}{yellow!20}
```

`\bggradient{blue!20}{red!20}`

Notice that this slide has no footer. This is accomplished by marking the frame as 'plain'.

```
(\bggradient{blue!15}{orange!15})
```

`\bggradient{purple!15}{orange!15}`

`\bggradient{blue!15}{green!15}`

\bggradient{green!15}{yellow!15}

$$I = -\frac{1}{16\pi G} \int_{\mathcal{M}} d^{n+1}x \sqrt{g} (R - 2\Lambda) - \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} K$$

$$- \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} \left(\frac{n-1}{\ell^2} + \frac{\ell}{2(n-2)} \mathcal{R} + \dots \right)$$

$$I = -\frac{1}{16\pi G} \int_{\mathcal{M}} d^{n+1}x \sqrt{g} (R - 2\Lambda) - \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} K$$

$$- \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} \left(\frac{n-1}{\ell^2} + \frac{\ell}{2(n-2)} \mathcal{R} + \dots \right)$$

$$I = -\frac{1}{16\pi G} \int_{\mathcal{M}} d^{n+1}x \sqrt{g} (R - 2\Lambda) - \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} K$$

$$- \frac{1}{8\pi G} \int_{\partial\mathcal{M}} d^n x \sqrt{\gamma} \left(\frac{n-1}{\ell^2} + \frac{\ell}{2(n-2)} \mathcal{R} + \dots \right)$$

Gradients with Three Colors

The command ‘bggradient’ produces a background gradient with three colors. It takes three arguments: the top, middle, and bottoms colors.

```
\bggradient{<top color>}{<middle color>}{<bottom color>}
```

For example, this page was produced with

```
\bggradient{gray3!60}{gray1!20}{gray3!60}
```


\bgggradient{skyblue3!40}{skyblue1!20}{skyblue3!40}

`\bggradient{skyblue3!80}{skyblue1!60}{skyblue3!80}`

Always test your presentation on a projector, to see what the colors look like. Sometimes they appear faint compared to how they look on the screen.

Outline

Introduction

First Section

Final Remarks

Macros for Boxes

Backgrounds

Custom Colors

Custom Colors

These are the custom colors defined in the Simple beamer style.

butter1	butter2	butter3
orange1	orange2	orange3
chocolate1	chocolate2	chocolate3
scarlet1	scarlet2	scarlet3
plum1	plum2	plum3
skyblue1	skyblue2	skyblue3
chameleon1	chameleon2	chameleon3

You can make them lighter by appending `!n` to the name, where n is a number between 0 and 100. I am using $n = 20$ here.

butter1

butter2

butter3

orange1

orange2

orange3

chocolate1

chocolate2

chocolate3

scarlet1

scarlet2

scarlet3

plum1

plum2

plum3

skyblue1

skyblue2

skyblue3

chameleon1

chameleon2

chameleon3

The custom colors were taken from the Tango Project.
<http://tango.freedesktop.org/>